

KAP1/TIF1 $\beta$  mouse mAb

<b>Catalog No :</b>	YM1235
<b>Reactivity :</b>	Human
<b>Applications :</b>	WB;ICC;IHC;IP
<b>Target :</b>	TIF1 $\beta$
<b>Gene Name :</b>	trim28
<b>Human Gene Id :</b>	10155
<b>Human Swiss Prot No :</b>	Q13263
<b>Mouse Swiss Prot No :</b>	Q62318
<b>Immunogen :</b>	Purified recombinant human KAP1 / TIF1 beta protein fragments expressed in E.coli.
<b>Specificity :</b>	This antibody detects endogenous levels of KAP1 / TIF1 beta and does not cross-react with related proteins.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	wb 1:1000 icc 1:100
<b>Purification :</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	110kD
<b>Background :</b>	The protein encoded by this gene mediates transcriptional control by interaction

with the Kruppel-associated box repression domain found in many transcription factors. The protein localizes to the nucleus and is thought to associate with specific chromatin regions. The protein is a member of the tripartite motif family. This tripartite motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. [provided by RefSeq, Jul 2008],

**Function :**

domain:Contains one Pro-Xaa-Val-Xaa-Leu (PxVxL) motif, which is required for interaction with chromoshadow domains. This motif requires additional residues -7, -6, +4 and +5 of the central Val which contact the chromoshadow domain.,domain:The HP1 box is both necessary and sufficient for HP1 binding. The RING finger domain and the B-box domains mediate interaction with CEBPB. The PHD domain enhances the CEBPB transcriptional activity.,function:Forms a complex with a KRAB-domain transcription factor and increases the efficiency of KRAB-mediated repression. Silences transcription through an interaction with HP1 proteins. Acts as a corepressor of transcription for the KRAB zinc finger proteins and as a moderator of the repression activity. May play a role as a coactivator for CEBPB and NR3C1 in the transcriptional activation of the Alpha-1-acid glycoprotein gene.,PTM:Phosphorylated upon DNA


**Subcellular Location :**

Nucleus . Associated with centromeric heterochromatin during cell differentiation through CBX1. .


**Expression :**

Expressed in all tissues tested including spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes.


## Products Images




Western blot detection of KAP1 / TIF1 beta in 293T and HepG2 cell lysates using KAP1 / TIF1 beta mouse mAb (1:1000 diluted). Observed band size: 110KDa.




Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-KAP1 / TIF1 beta mouse mAb (dilution 1:100).



Immunoprecipitation analysis of Hela cell lysates using KAP1 / TIF1 beta mouse mAb.



IHC of paraffin-embedded human Spleen using anti-KAP1 / TIF1 beta diluted 1/500-1/1000.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).