

## 2A5B Polyclonal Antibody

<b>Catalog No :</b>	YN1277
<b>Reactivity :</b>	Human;Rat;Mouse;
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	2A5B
<b>Fields :</b>	>>mRNA surveillance pathway;>>Sphingolipid signaling pathway;>>Oocyte meiosis;>>PI3K-Akt signaling pathway;>>AMPK signaling pathway;>>Adrenergic signaling in cardiomyocytes;>>Dopaminergic synapse;>>Human papillomavirus infection
<b>Gene Name :</b>	PPP2R5B
<b>Protein Name :</b>	Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit beta isoform (PP2A B subunit isoform B'-beta) (PP2A B subunit isoform B56-beta) (PP2A B subunit isoform PR61-beta) (PP2A B subunit iso
<b>Human Gene Id :</b>	5526
<b>Human Swiss Prot No :</b>	Q15173
<b>Immunogen :</b>	Synthesized peptide derived from human protein . at AA range: 160-240
<b>Specificity :</b>	2A5B Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml

<b>Storage Stability :</b>	<u>-15 °C to -25 °C/1 year(Do not lower than -25 °C)</u>
<b>Observed Band :</b>	<u>54kD</u>
<b>Cell Pathway :</b>	<u>Oocyte meiosis;WNT;WNT-T CELL</u>
<b>Background :</b>	<u>The product of this gene belongs to the phosphatase 2A regulatory subunit B family. Protein phosphatase 2A is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The B regulatory subunit might modulate substrate selectivity and catalytic activity. This gene encodes a beta isoform of the regulatory subunit B56 subfamily. [provided by RefSeq, Jul 2008],</u>
<b>Function :</b>	<u>function:The B regulatory subunit might modulate substrate selectivity and catalytic activity, and also might direct the localization of the catalytic enzyme to a particular subcellular compartment.,induction:By retinoic acid; in neuroblastoma cell lines.,PTM:At least isoform Beta-1 is phosphorylated on serine residues.,similarity:Belongs to the phosphatase 2A regulatory subunit B56 family.,subunit:PP2A consists of a common heterodimeric core enzyme, composed of a 36 kDa catalytic subunit (subunit C) and a 65 kDa constant regulatory subunit (PR65 or subunit A), that associates with a variety of regulatory subunits. Proteins that associate with the core dimer include three families of regulatory subunits B (the R2/B/PR55/B55, R3/B"/PR72/PR130/PR59 and R5/B'/B56 families), the 48 kDa variable regulatory subunit, viral proteins, and cell signaling molecules. Interacts with SGOL1.,tissue sp</u>
<b>Subcellular Location :</b>	<u>Cytoplasm .</u>
<b>Expression :</b>	<u>Highest expression in brain.</u>

## Products Images