

Recombinant Human IL1B protein ,C- His Tag

YD2203 Catalog No:

Reactivity: Human

>90% as determined by SDS-PAGE **Purity:**

P01584

In Vivo: Please contact with the lab for this information.

Sequence: A DNA sequence encoding the human IL1B(Ala117~Ser269) was fused with the

C-terminal His Tag

Human Swiss Prot

No:

Formulation: Supplied as solution form in PBS pH7.5./ Lyophilized from PBS pH7.5.

Source: E.coli

Dilution: Reconstitute in sterile water for a stock solution

Concentration: >90% as determined by SDS-PAGE

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 **Storage Stability:**

to 8 °C for one week. Store at -20 to -80 °C for twelve months from the date of

receipt.

Subcellular

Cytoplasm, cytosol . Secreted . Lysosome . Secreted, extracellular exosome . The precursor is cytosolic (PubMed:15192144). In response to inflammasome-Location: activating signals, such as ATP for NLRP3 inflammasome or bacterial flagellin for

NLRC4 inflammasome, cleaved and secreted (PubMed:24201029,

PubMed:33377178, PubMed:33883744). Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744). In contrast, the precursor form is not released, due to the presence of an acidic region that is proteolytically removed by CASP1 during maturation (PubMed:33883744). The secretion is dependent on protein unfolding

and facilitated by the cargo receptor TMED10 (PubMed:32272059)...

Expression: Expressed in activated monocytes/macrophages (at protein level).

1/2



Products Images