

LSD1 rabbit-FC recombinant protein

Catalog No :	YD3141
Reactivity :	Human;
Purity :	>90% as determined by SDS-PAGE
Gene Name :	KDM1/LSD1
Protein Name :	Lysine-specific histone demethylase 1A
Sequence :	Amino acid:21-210,with rabbit FC tag.
Human Gene Id :	23028
Human Swiss Prot No :	O60341
Formulation :	Phosphate-buffered solution
Source :	Mammalian cells
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Background :	<p>This gene encodes a nuclear protein containing a SWIRM domain, a FAD-binding motif, and an amine oxidase domain. This protein is a component of several histone deacetylase complexes, though it silences genes by functioning as a histone demethylase. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2009],</p>
Function :	<p>cofactor:FAD.,domain:The SWIRM domain may act as an anchor site for a histone tail.,function:Histone demethylase that demethylates 'Lys-4' of histone H3, a specific tag for epigenetic transcriptional activation, thereby acting as a corepressor. Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed. Demethylates both mono- and di-methylated 'Lys-4' of histone H3. May play a role in the repression of neuronal genes. Alone, it is unable to demethylate H3 'Lys-4' on nucleosomes and requires the presence of RCOR1/CoREST to achieve such activity. May also demethylate 'Lys-9' of histone H3, a specific tag for epigenetic transcriptional repression, thereby leading to derepression of androgen receptor target genes. Demethylates di-methylated 'Lys-370' of p53/TP53 which prevents interaction of p53/TP53 with TP53BP1 and represses p53/TP53-mediate</p>

SubcellularNucleus .

Location :**Expression :**Ubiquitously expressed.

Products Images