

## **CTGF Polyclonal Antibody**

Catalog No: YT6207

**Reactivity:** Human; Mouse; Rat

**Applications:** IHC;IF;WB

Target: CTGF

**Fields:** >>Apelin signaling pathway;>>Hippo signaling pathway

Gene Name: CTGF CCN2 HCS24 IGFBP8

P29279

Protein Name: CTGF

Human Gene Id: 1490

**Human Swiss Prot** 

No:

**Immunogen:** Synthesized peptide derived from human CTGF AA range: 130-210

**Specificity:** This antibody detects endogenous levels of human CTGF

**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, lgG

**Dilution :** IHC 1:50-200, WB 1:500-2000. IF 1:50-200

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Concentration**: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band:** 37kD



**Background:** The protein encoded by this gene is a mitogen that is secreted by vascular

endothelial cells. The encoded protein plays a role in chondrocyte proliferation and differentiation, cell adhesion in many cell types, and is related to platelet-derived growth factor. Certain polymorphisms in this gene have been linked with a

higher incidence of systemic sclerosis. [provided by RefSeq, Nov 2009],

**Function:** function:Major connective tissue mitoattractant secreted by vascular endothelial

cells. Promotes proliferation and differentiation of chondrocytes. Mediates

heparin- and divalent cation-dependent cell adhesion in many cell types including fibroblasts, myofibroblasts, endothelial and epithelial cells. Enhances fibroblast

growth factor-induced DNA synthesis., similarity: Belongs to the CCN family., similarity: Contains 1 CTCK (C-terminal cystine knot-like)

domain.,similarity:Contains 1 IGFBP N-terminal domain.,similarity:Contains 1 TSP type-1 domain.,similarity:Contains 1 VWFC domain.,subunit:Monomer.,

Subcellular Location:

Secreted, extracellular space, extracellular matrix. Secreted.

**Expression:** Expressed in bone marrow and thymic cells. Also expressed one of two Wilms

tumors tested.

## **Products Images**

2/2