

LC3B Polyclonal Antibody

Catalog No :	YN5524
Reactivity :	Human;Mouse;Rat
Applications :	WB;IF;IHC
Target :	LC3B
Fields :	>>Mitophagy - animal;>>Autophagy - animal;>>Ferroptosis;>>Apelin signaling pathway;>>NOD-like receptor signaling pathway;>>Amyotrophic lateral sclerosis;>>Pathways of neurodegeneration - multiple diseases;>>Shigellosis;>>Kaposi sarcoma-associated herpesvirus infection
Gene Name :	MAP1LC3B
Protein Name :	Microtubule-associated proteins 1A/1B light chain 3B
Human Gene Id :	81631
Human Swiss Prot No :	Q9GZQ8
Mouse Swiss Prot No :	Q9CQV6
Rat Swiss Prot No :	Q62625
Immunogen :	Recombinant Protein of MAP LC3 β
Specificity :	The antibody detects endogenous MAP LC3 β protein.
Formulation :	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:1000-2000 IHC 1:200-500 IF 1:200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Storage Stability : -15°C to -25°C/1 year (Do not lower than -25°C)

Observed Band : 14,16kD

Background : The product of this gene is a subunit of neuronal microtubule-associated MAP1A and MAP1B proteins, which are involved in microtubule assembly and important for neurogenesis. Studies on the rat homolog implicate a role for this gene in autophagy, a process that involves the bulk degradation of cytoplasmic component. [provided by RefSeq, Jul 2008],

Function : caution:PubMed:12740394 has shown that the protein is cleaved at Lys-122 but PubMed:15355958 has shown that the cleavage site is at Gly-120 as in other mammalian orthologs.,function:Probably involved in formation of autophagosomal vacuoles (autophagosomes).,PTM:The precursor molecule is cleaved by APG4B/ATG4B to form LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form LC3-II.,similarity:Belongs to the MAP1 LC3 family.,subcellular location:LC3-II binds to the autophagic membranes.,subunit:3 different light chains, LC1, LC2 and LC3, can associate with MAP1A and MAP1B proteins.,tissue specificity:Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver.,

Subcellular Location : Cytoplasmic vesicle, autophagosome membrane ; Lipid-anchor . Endomembrane system ; Lipid-anchor . Mitochondrion membrane ; Lipid-anchor . Cytoplasm, cytoskeleton . Cytoplasmic vesicle . LC3-II binds to the autophagic membranes. LC3-II localizes with the mitochondrial inner membrane during Parkin-mediated mitophagy (PubMed:28017329). Localizes also to discrete punctae along the ciliary axoneme. .

Expression : Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver.

Products Images