

FGFP1 Polyclonal Antibody

Catalog No :	YN1743
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	FGFP1
Gene Name :	FGFBP1 FGFBP HBP17
Protein Name :	Fibroblast growth factor-binding protein 1 (FGF-BP) (FGF-BP1) (FGF-binding protein 1) (FGFBP-1) (17 kDa heparin-binding growth factor-binding protein) (17 kDa HBGF-binding protein) (HBp17)
Human Gene Id :	9982
Human Swiss Prot No :	Q14512
Mouse Swiss Prot No :	O70514
Rat Swiss Prot No :	Q9QY10
Immunogen :	Synthesized peptide derived from human protein . at AA range: 110-190
Specificity :	FGFP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 25kD


Background : This gene encodes a secreted fibroblast growth factor carrier protein. The encoded protein plays a critical role in cell proliferation, differentiation and migration by binding to fibroblast growth factors and potentiating their biological effects on target cells. The encoded protein may also play a role in tumor growth as an angiogenic switch molecule, and expression of this gene has been associated with several types of cancer including pancreatic and colorectal adenocarcinoma. A pseudogene of this gene is also located on the short arm of chromosome 4. [provided by RefSeq, Nov 2011],

Function : function:Acts as a carrier protein that release fibroblast-binding factors (FGFs) from the extracellular matrix (EM) storage and thus enhance the mitogenic activity of FGFs. Enhances FGF2 signaling during tissue repair, angiogenesis and in tumor growth.,induction:Up-regulated in epithelial cells after skin injury. Keratinocyte mitogens.,miscellaneous:Expression is significantly up-regulated in carcinogen-induced skin tumors, various squamous cell carcinomas, some colon cancer cell lines and tumors.,similarity:Belongs to the fibroblast growth factor-binding protein family.,subcellular location:Extracellular and plasma membrane-associated. Colocalizes with HSPG2 in the pericellular environment of squamous cell carcinomas.,subunit:Found in a complex with FGFBP1, FGF1 and FGF2. Interacts with FGF1, FGF2, FGF7, FGF10, FGF22 and HSPG2.,tissue specificity:Expressed in the suprabasal region of t

Subcellular Location : Secreted, extracellular space . Cell membrane ; Peripheral membrane protein . Extracellular and plasma membrane-associated. Colocalizes with HSPG2 in the pericellular environment of squamous cell carcinomas. .

Expression : Expressed in the suprabasal region of the epidermis, in hair follicles, the basement membrane at the dermo-epidermal junction (occasionally extending into the basement membrane of dermal blood vessels), wounded skin and several invasive squamous cell carcinomas (at protein level). Expressed in normal and wounded skin and various squamous cell carcinomas.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night