

β Actin (PT0519R) PT® Rabbit mAb

Catalog No: YM8343

Reactivity: Human; Mouse; Rat;

Applications: WB;IHC;IF;IP;ELISA

Target: Actin β

Fields: >>Rap1 signaling pathway;>>Phagosome;>>Apoptosis;>>Hippo signaling

pathway;>>Focal adhesion;>>Adherens junction;>>Tight junction;>>Platelet activation;>>Neutrophil extracellular trap formation;>>Leukocyte transendothelial migration;>>Thermogenesis;>>Regulation of actin cytoskeleton;>>Thyroid hormone signaling pathway;>>Oxytocin signaling pathway;>>Gastric acid secretion;>>Amyotrophic lateral sclerosis;>>Bacterial invasion of epithelial

cells;>>Vibrio cholerae infection;>>Pathogenic Escherichia coli

infection;>>Shigellosis;>>Salmonella infection;>>Yersinia infection;>>Influenza A;>>Proteoglycans in cancer;>>Hepatocellular carcinoma;>>Hypertrophic cardiomyopathy;>>Arrhythmogenic right ventricular cardiomyopathy;>>Dilated cardiomyopathy;>>Viral myocarditis;>>Fluid shear stress and atherosclerosis

Gene Name: ACTB

Protein Name: Actin cytoplasmic 1

Human Gene Id: 60

Human Swiss Prot

No:

Mouse Gene ld: 11461

Mouse Swiss Prot

No:

P60710

P60709

Rat Gene Id: 81822

Rat Swiss Prot No: P60711

Specificity: endogenous

Formulation: PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA



Source : Monoclonal, rabbit, IgG, Kappa

Dilution: IHC 1:2000-1:20000;WB 1:2000-1:10000;IF 1:200-1:1000;ELISA

1:5000-1:20000;IP 1:50-1:200;

Purification: Protein A

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 42kD

Observed Band: 42kD

Cell Pathway: Focal adhesion; Adherens_Junction; Adherens_Junction; Leukocyte

transendothelial migration; Regulates Actin and Cytoskeleton; Vibrio cholerae

infection;Pathogenic Escherichia coli infection;Hypertrophic ca

Background: This gene encodes one of six different actin proteins. Actins are highly

conserved proteins that are involved in cell motility, structure, and integrity. This actin is a major constituent of the contractile apparatus and one of the two

nonmuscle cytoskeletal actins. [provided by RefSeg, Jul 2008],

Function : disease:Defects in ACTB are a cause of dystonia juvenile-onset (DYTJ)

[MIM:607371]. DYTJ is a form of dystonia with juvenile onset. Dystonia is defined by the presence of sustained involuntary muscle contraction, often leading to abnormal postures. DYTJ patients manifest progressive, generalized, dopa-unresponsive dystonia, developmental malformations and sensory hearing loss., function: Actins are highly conserved proteins that are involved in various

types of cell motility and are ubiquitously expressed in all eukaryotic

cells.,miscellaneous:In vertebrates 3 main groups of actin isoforms, alpha, beta and gamma have been identified. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. The beta and gamma actins coexist in most cell types as components of the cytoskeleton and as

mediators of internal cell motility., similarity: Belongs to the


Subcellular Location:

Cytoplasm

Expression: B-cell lymphoma, Brain, Cajal-Retzius cell, Eye, Fetal brain

cortex, Foreskin, Hepatocellular car

Products Images



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti β Actin (PT0519R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: 3T3-L1 Lane 2: Hela Lane 3: C6 Predicted band size: 42kDa Observed band size: 42kDa