

## PP2BB Polyclonal Antibody

<b>Catalog No :</b>	YN2056
<b>Reactivity :</b>	Human;Rat;Mouse
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	PP2BB
<b>Fields :</b>	>>MAPK signaling pathway;>>Calcium signaling pathway;>>cGMP-PKG signaling pathway;>>Oocyte meiosis;>>Cellular senescence;>>Wnt signaling pathway;>>Axon guidance;>>VEGF signaling pathway;>>Osteoclast differentiation;>>C-type lectin receptor signaling pathway;>>Natural killer cell mediated cytotoxicity;>>Th1 and Th2 cell differentiation;>>Th17 cell differentiation;>>T cell receptor signaling pathway;>>B cell receptor signaling pathway;>>Long-term potentiation;>>Glutamatergic synapse;>>Dopaminergic synapse;>>Oxytocin signaling pathway;>>Glucagon signaling pathway;>>Renin secretion;>>Alzheimer disease;>>Amyotrophic lateral sclerosis;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Amphetamine addiction;>>Tuberculosis;>>Human cytomegalovirus infection;>>Human T-cell leukemia virus 1 infection;>>Kaposi sarcoma-associated herpesvirus infection;>>Human immunodeficiency virus 1 infection;>>PD-L1 expression and PD-1 checkpoint pathway in cancer;>>Lipid and atherosclerosis
<b>Gene Name :</b>	PPP3CB CALNA2 CALNB CNA2
<b>Protein Name :</b>	Serine/threonine-protein phosphatase 2B catalytic subunit beta isoform (EC 3.1.3.16) (CAM-PRP catalytic subunit) (Calmodulin-dependent calcineurin A subunit beta isoform)
<b>Human Gene Id :</b>	5532
<b>Human Swiss Prot No :</b>	P16298
<b>Mouse Swiss Prot No :</b>	P48453
<b>Rat Swiss Prot No :</b>	P20651
<b>Immunogen :</b>	Synthesized peptide derived from part region of human protein
<b>Specificity :</b>	PP2BB Polyclonal Antibody detects endogenous levels of protein.

<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	57kD
<b>Cell Pathway :</b>	MAPK_ERK_Growth;MAPK_G_Protein;Calcium;Oocyte meiosis;Apoptosis_Inhibition;Apoptosis_Mitochondrial;Apoptosis_Overview;WNT;WNT-T CELLAxon guidance;VEGF;Natural killer cell mediated cytotoxicity;T_Cell_
<b>Background :</b>	alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events,catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 1 Fe(3+) ion per subunit.,cofactor:Binds 1 zinc ion per subunit.,function:Calcium-dependent, calmodulin-stimulated protein phosphatase. This subunit may have a role in the calmodulin activation of calcineurin.,similarity:Belongs to the PPP phosphatase family.,similarity:Belongs to the PPP phosphatase family. PP-2B subfamily.,subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity.,
<b>Function :</b>	alternative products:Additional isoforms seem to exist. Calcineurin A beta isoform consists of at least two isoenzymes that may result from alternative splicing events,catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 1 Fe(3+) ion per subunit.,cofactor:Binds 1 zinc ion per subunit.,function:Calcium-dependent, calmodulin-stimulated protein phosphatase. This subunit may have a role in the calmodulin activation of calcineurin.,similarity:Belongs to the PPP phosphatase family.,similarity:Belongs to the PPP phosphatase family. PP-2B subfamily.,subunit:Composed of two components (A and B), the A component is the catalytic subunit and the B component confers calcium sensitivity.,
<b>Subcellular Location :</b>	Cytoplasm .
<b>Expression :</b>	Teratocarcinoma,Testis,

## Products Images