

STAM1 Polyclonal Antibody

Catalog No :	YN2826
Reactivity :	Human;Mouse
Applications :	WB;ELISA
Target :	STAM1
Fields :	>>Endocytosis;>>JAK-STAT signaling pathway
Gene Name :	STAM STAM1
Protein Name :	Signal transducing adapter molecule 1 (STAM-1)
Human Gene Id :	8027
Human Swiss Prot No :	Q92783
Mouse Swiss Prot No :	P70297
Immunogen :	Synthesized peptide derived from part region of human protein AA range: 156-206
Specificity :	STAM1 Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 59kD

Cell Pathway : Endocytosis;Jak_STAT;

Background : This gene encodes a member of the signal-transducing adaptor molecule family. These proteins mediate downstream signaling of cytokine receptors and also play a role in ER to Golgi trafficking by interacting with the coat protein II complex. The encoded protein also associates with hepatocyte growth factor-regulated substrate to form the endosomal sorting complex required for transport-0 (ESCRT-0), which sorts ubiquitinated membrane proteins to the ESCRT-1 complex for lysosomal degradation. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Feb 2011],

Function : function:Involved in intracellular signal transduction mediated by cytokines and growth factors. Upon IL-2 and GM-CSL stimulation, it plays a role in signaling leading to DNA synthesis and c-myc induction. May also play a role in T-cell development. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with HGS (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes.,PTM:Phosphorylated on Tyr-198. Phosphorylation occurs in response to IL-2, IL-3, IL-4, IL-7, GM-CSF, EGF and PDGF.,similarity:Belongs to the STAM family.,similarity:Contains 1 ITAM domain.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UIM (ubiquitin-interacting motif) repeat.,similarity:Contains 1 VHS domain.,subunit:Com

Subcellular Location : Cytoplasm . Early endosome membrane ; Peripheral membrane protein ; Cytoplasmic side .

Expression : Ubiquitously expressed.

Products Images