ApplicationsWB,IHC,IF,ELISA



TNF-α Rabbit pAb

CatalogNo: YT4689 Orthogonal Validated 💽

Key Features

Host Species Reactivity

Rabbit
 Human, Mouse, Rat

MW Isotype
• 16kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-1:2000 IHC: 1:100-300 ELISA 1:20000 IF 1:100-300

Not yet tested in other applications.

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human TNFA. AA

range:141-190

Specificity TNF- α Polyclonal Antibody detects endogenous levels of TNF- α protein.

Target Information

Gene name

TNF TNFA TNFSF2

Protein Name

Tumor necrosis factor (Cachectin) (TNF-alpha) (Tumor necrosis factor ligand superfamily member 2) (TNF-a)

Organism	Gene ID	UniProt ID
Human	<u>7124;</u>	<u>P01375;</u>
Mouse	<u>21926;</u>	<u>P06804;</u>
Rat	<u>24835;</u>	<u>P16599;</u>

Cellular Localization

Cell membrane; Single-pass type II membrane protein.; [Tumor necrosis factor, membrane form]: Membrane; Single-pass type II membrane protein.; [Tumor necrosis factor, soluble form]: Secreted .; [C-domain 1]: Secreted.; [C-domain 2]: Secreted.

Tissue specificity Blood, Peripheral blood leukocyte, Placenta, Prostatic carcinoma,

Function

Disease: Cachexia accompanies a variety of diseases, including cancer and infection, and is characterized by general ill health and malnutrition., Disease: Genetic variations in TNF are associated with susceptibility to hepatitis B virus infection (HBV infection) [MIM:610424]. Approximately one third of all cases of cirrhosis and half of all cases of hepatocellular carcinoma can be attributed to chronic HBV infection. HBV infection may result in subclinical or asymptomatic infection, acute self-limited hepatitis, or fulminant hepatitis requiring liver transplantation., Disease: Genetic variations in TNF are associated with susceptibility to psoriatic arthritis [MIM:607507]. Psoriasis is a chronic inflammatory dermatosis that affects approximately 2% of the population. It is characterized by red, scaly skin lesions that are usually found on the scalp, elbows, and knees, and may be associated with severe arthritis. Psoriatic arthritis has been defined as an inflammatory arthritis usually without any rheumatoid factor in serum (seronegative arthritis) associated with psoriasis., Function: Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation., online information: The Singapore human mutation and polymorphism database, online information: Tumor necrosis factor alpha entry,PTM:O-glycosylated; glycans contain galactose, N-acetylgalactosamine and Nacetylneuraminic acid.,PTM:The membrane form, but not the soluble form, is phosphorylated on serine residues. Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1.,PTM: The soluble form derives from the membrane form by proteolytic processing., similarity: Belongs to the tumor necrosis factor family., subunit: Homotrimer.,

I Validation Data

Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **TNF-α Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents